

Higher Entropy

What is "Higher Entropy?"

How much ignorance do I have? \rightsquigarrow What is it that I don't know?

(relative) Entropy \rightsquigarrow "Linear Space" of Random Variables

E.g. $\hat{\mu} : \Omega \rightarrow \mathbb{R}_{\geq 0}$

$$S(\hat{\mu}) = \sum_{\omega \in \Omega} \hat{\mu}_\omega \log \hat{\mu}_\omega$$

$\rightsquigarrow L^2(\Omega, \mu \neq 0)$

$q \in \mathbb{H} \mid \operatorname{Re} q \geq 1, \mathbb{C}$

measuring these resolves ignorance

$\cdot \|f\|_2 = \left| \sum_{\omega} (f_{\omega}^* f_{\omega}) \right|^{1/2}$

Homotopical Perspective: Study multipartite Systems?
 e.g. $M_{AB} : \Omega_A \times \Omega_B \rightarrow \mathbb{R}_{\geq 0}$

How much info. is shared? \rightsquigarrow What info. is shared?

Mutual Information:

$$I(M_{AB}, \{A, B\}) = S_A + S_B - S_{AB}$$

\rightsquigarrow "Space" / (co-)simplicial obj.

$\hat{\mu}_A(\omega_A) = \sum_{\omega_B \in \Omega_B} \hat{\mu}(\omega_A, \omega_B)$

$M_\phi \leftarrow M_A \boxplus M_B \leftarrow M_{AB}$

Multipartite Measures

"Spaces" (Simplicial measures)

Homology

Graded Vector Spaces (+ extra data)

("Euler Char")

Euler Char

Mutual Information

Holomorphic Functions of q

$$\frac{d}{dq} \Big|_{q=1}$$

OF q

$q \rightarrow 0$


\mathbb{R}

\mathbb{Z}

$$\mu: \Omega_A \times \Omega_B \rightarrow \mathbb{R}_{\geq 0}$$


Probability Matrix (bipartite)

$$\begin{matrix}
 & a_1 & a_2 & \dots \\
 b_1 & \mu_{11} & & \\
 b_2 & 0 & \mu_{22} & \\
 \vdots & & & \mu_{33}
 \end{matrix}$$



$$\chi_\mu^{2q} = \sum_i \mu_{ii}^{AB q} - 1$$

State index $\in \mathcal{O}(\mathbb{C}_q \times \mathbb{C}_r \times \mathbb{C}_s \times \mathbb{C}_t)$



Tsallis/Rényi deformed MI $\in \mathcal{O}(\mathbb{C}_q \times \mathbb{C}_r)$

Rényi MI

Tsallis MI

MI

$$H^0 \cong \{ (\alpha, \beta) : \text{Cov}(\alpha, \beta) = |\text{Var}(\alpha)|^2 = |\text{Var}(\beta)|^2 \}$$

$$\alpha: \Omega_A \rightarrow \mathbb{C}$$

$$\beta: \Omega_B \rightarrow \mathbb{C}$$

Why (Should this be possible)?

• Correlations among Subsystems \longleftrightarrow Obstruction to Factorizability
 ⚡ "Space/Cohomology"

• Mutual Info. looks like an EC

$$I_{AB} \stackrel{?}{=} \dim [M_A \oplus M_B \hookrightarrow M_{AB}]$$

$$= \underbrace{\dim(M_A) + \dim(M_B)}_{\text{"S}_A?"} - \dim(M_{AB})$$

- μ Factorizes in any way $\Rightarrow MI = 0$
 \Leftarrow
- MI measures info shared among all subsystems

The Category of Measures \rightarrow States

Def: A state is a pair of a \mathbb{C} -alg. A and a positive linear map $\mu: A \rightarrow \mathbb{C}$

$$\mu(a^*a) \geq 0 \quad \forall a$$

Ex: $A = \prod_{i=1}^N M_{n_i} \mathbb{C}$, $\mu: A \rightarrow \mathbb{C}$
 $a \mapsto \sum_{i=1}^N \text{Tr}_i [\hat{\mu}_i a] \in (M_n \mathbb{C})_{\geq 0}$

- purely classical: $n_i = 1 \quad \forall i \Rightarrow \hat{\mu}_i \in \mathbb{R}_{\geq 0}$
- purely quantum: $N=1 \quad (n_1 > 0)$.

Fix A an algebra

Def: State_A has


• Objects $\mu: A \rightarrow \mathbb{C}$ states

• Morphisms: $\mu \xrightarrow{!} \nu$; $F: \nu \leq \mu$
 $(\nu(a^*a) \leq \mu(a^*a) \forall a)$

(RN deriv $\nu = F \cdot \mu$)

" $L(X, \mu)_{+ \leq}$ "

State is the (Grothendieck op-Fibration)



$$\begin{pmatrix} F_A: \mu_B \rightarrow \mu_A \\ \text{is given by} \\ F: A \rightarrow B \text{ s.t.} \\ F^* \mu_B \leq \mu_A \end{pmatrix}$$

(Isos are "unitary maps" $u: A \rightarrow B$ s.t. $u^* \mu_B = \mu_A$)
 \uparrow
 $^* \text{-isos}$

• State has Coproducts:

$$\begin{aligned} \mu_A \boxplus \mu_B &: A \times B \rightarrow \mathbb{C} \\ (a, b) &\mapsto \mu_A(a) + \mu_B(b) \end{aligned}$$

• State has \otimes

$$\begin{aligned} \mu_A \otimes \mu_B &: A \otimes B \rightarrow \mathbb{C} \\ a \otimes b &\mapsto \mu_A(a) \mu_B(b) \end{aligned}$$

Dimension

\mathcal{C} a category w/ Coproducts and \otimes :

$$\dim : \{ \text{Iso classes of } \mathcal{C} \} \longrightarrow \mathbb{R} \leftarrow \text{ring}$$

$$\dim (C_1 \perp C_2) = \dim (C_1) + \dim (C_2)$$

$$\dim (C_1 \otimes C_2) = \dim (C_1) \dim (C_2)$$

Ex:

\mathcal{C}	dim
Set	Card
Vect $_{\mathbb{R}}$	dim $_{\mathbb{R}}$
$(V, f: V \rightarrow V)$	Tr(f^n), $n \in \mathbb{Z}_{\geq 0}$
Graded V.S. $V \oplus_k V^k$	$\sum_k (-1)^n \dim_{\mathbb{R}} (V^k)$

$\mathcal{C} = \text{Fin Meas} \leftarrow \text{States on } \mathbb{C}^n / \hat{\mu}: \Omega \rightarrow \mathbb{C}, \mathbb{R} = \mathcal{O}(\mathbb{C})$

$$\dim_q (M) = \sum_{\hat{\mu}_i} \hat{\mu}_i^q$$

$\mathcal{C} = \text{Fin State}, \mathbb{R} = \mathcal{O}(\mathbb{C}^3)$

$$\dim_{\alpha, \mathbb{Z}, \mathbb{R}} (M) = \sum_i n_i^\alpha \text{Tr} [\hat{\mu}_i^q]$$

(EC of $\mu \leftarrow M$)

This inspires

$$S_{\mathbb{Z}, \mathbb{R}}^{\text{TR}} (M) = \frac{1}{q-1} [1 - \dim_{\alpha, \mathbb{Z}, \mathbb{R}} (M)]$$

$q \rightarrow 0$ limit: $\dim \in \mathbb{Z}$

Classical case: $\dim = \# \text{ points w/ } M \neq 0$ "L⁰ norm"

Thus, we expect an association:

State \rightsquigarrow Vector Space (or Set)

GNS Modules

- There is a Functor:

$$\text{GNS} : \text{State}_A \longrightarrow_A \text{Mod}$$

$$\mu \longmapsto A/\mathcal{I}_\mu, \quad \mathcal{I}_\mu = \{a : \mu(a^*a) = 0\}$$

$$(\mu \longrightarrow \nu) \longmapsto \mathcal{Q} : A/\mathcal{I}_\mu \longrightarrow A/\mathcal{I}_\nu \leftarrow \begin{array}{l} \text{"quotient"} \\ \text{map} \\ \text{"RN-deriv"} \end{array}$$


- $\text{GNS}(\hat{\mu} : \Omega \rightarrow \mathbb{R}_{\geq 0}) \cong \mathbb{C}[\Omega_{\mu \neq 0}]$

- q Completions:

$$\begin{aligned} \|\cdot\|_q : \text{GNS}(\mu) &\longrightarrow \mathbb{R}_{\geq 0} \\ [a] &\longmapsto \left| \mu((a^*a)^{q/2}) \right|^{1/\text{Re } q} \end{aligned}$$

$\rightsquigarrow L^q$ Spaces

Further Work :



- G-equivariant entropy
- Link invariants
- Categorify Weir inequalities for entropy / Modular Flow
- Entropy for Finite Fields
- Emergent Geometry / Spacetime